

8-23-2013

Epidemiology News

Georgia Southern University

Follow this and additional works at: <https://digitalcommons.georgiasouthern.edu/epid-news-online>

Part of the [Epidemiology Commons](#)

Recommended Citation

Georgia Southern University, "Epidemiology News" (2013). *Epidemiology News (through 6/2018)*. 10.
<https://digitalcommons.georgiasouthern.edu/epid-news-online/10>

This article is brought to you for free and open access by the Biostatistics, Epidemiology, and Environmental Health Sciences Department Publications at Digital Commons@Georgia Southern. It has been accepted for inclusion in Epidemiology News (through 6/2018) by an authorized administrator of Digital Commons@Georgia Southern. For more information, please contact digitalcommons@georgiasouthern.edu.

Perceived Barriers

August 23, 2013



A collaborative study including Dr. Yelena Tarasenko at Jiann-Ping Hsu College of Public Health Georgia Southern University examines The Relationship Between Perceived Burden of Chronic Conditions and Colorectal Cancer Screening Among Appalachian Residents. As the population living with several concurrent chronic conditions or multiple morbidity (MM) increases, understanding how to effectively fit prevention efforts into disease management becomes more important, particularly among rural, underserved populations. Compared to their urban counterparts, rural residents suffer higher rates of disease, receive fewer preventive services, and often live in environments limiting access to optimal medical care. This study describes rural residents' perceived burdens of disease management and explores the relationship between these burdens, as proxies of individuals' competing demands, and colorectal cancer screening (CRCS).

We conducted a cross-sectional study, based on telephone survey data from 1,012 Appalachian residents, ages 50-75, with 1 or more chronic conditions. Measures of perceived MM burdens were developed based on 85 pilot interviews previously undertaken with providers and patients with MM residing in Appalachian Kentucky. Many participants (81%) agreed with 1 or more statements indicating perceived burdens of disease management effects on receiving CRCS. A higher percentage of rural (vs non-rural) Appalachians perceived burdens related to physician's recommendation, preparation to colonoscopy, and time management and affordability of both current diseases and screening. These differences did not modify the overall association between perceiving MM as burdensome and forgoing CRCS. The negative effect on CRCS of perceived burdens related to interaction with physician and time management was lower for participants with multiple rather than single morbidity. Future research designed to address perceived burdens of MM and improved interaction with health care providers may enhance critical prevention efforts among vulnerable populations.

Share: [f](#) [t](#) [g+](#) [+](#)

Posted in [Archive](#), [Epidemiology](#), [Health Policy & Mgmt](#), [JPHCOPH](#), [Research](#)