Summer 2017

Hepatitis C and HIV Screening

Follow this and additional works at: <https://digitalcommons.georgiasouthern.edu/jgpha>

Part of the [Public Health Commons](#)

Recommended Citation

Perry, Merry L. (2017) "Hepatitis C and HIV Screening," *Journal of the Georgia Public Health Association*: Vol. 7 : No. 1 , Article 48.

DOI: 10.21633/jgpha.7.149

Available at: <https://digitalcommons.georgiasouthern.edu/jgpha/vol7/iss1/48>

This conference abstract is brought to you for free and open access by the Journals at Digital Commons@Georgia Southern. It has been accepted for inclusion in Journal of the Georgia Public Health Association by an authorized administrator of Digital Commons@Georgia Southern. For more information, please contact digitalcommons@georgiasouthern.edu.

Conference Abstract

Hepatitis C and HIV screening

Merry L. Perry, DNP, ANP-BC

Doctor of Nursing Practice

Corresponding author: Merry L. Perry • Doctor of Nursing Practice • 106 East Broad Street, Savannah, GA 31401 • 912-527-2727 • mperry@cvcphc.com

Background: Chronic hepatitis C virus (HCV) affects more than 3 million people in the United States. Current HCV treatment offers a shorter treatment interval with fewer adverse reactions as well as improved cure rates of 96%. The increasing rates of hepatitis C infections in adults have been accompanied by multifaceted adverse health outcomes. The purpose of this screening program is to identify HCV-infected individuals, which will lead to clinical interventions and treatment that will improve health outcomes compared to no screening.

Methods: The Centers for Disease Control and Prevention guidelines recommend that anyone with risk factors should be screened for HCV infection. In addition, all persons born between 1945 and 1965 should be screened once in their lifetime. Screening at Curtis V. Cooper Primary HealthCare Inc. (CVCPHC) includes individuals 18 and over for HCV and HIV.

Results: CVCPHC's screening program was implemented in January 2016. As of October 1, 2016, 4054 individuals had been screened for HCV with 187 antibody-positive. Of the individuals with positive antibody tests, 179 were RNA tested, and of these, 122 tested positive. Seventy four percent of those with a positive RNA test were linked to care. Eighty five percent of individuals with a positive RNA test were born between 1945 and 1965. The current case reports consist of 57 active HCV patients on medication and/or have completed therapy; 10 have a viral load not detected (ND) > 4-6 weeks after starting medication; 15 completed therapy with a ND viral load resulting in a 100% cure rate; 12 are currently on medication < 4 weeks; 15 were referred to GI; and five deferred treatment.

Conclusions: Screening for HCV and HIV lead to the appropriate interventions and treatments for persons infected, preventing the progression of liver disease and reducing morbidity and possibly mortality.

Key words: hepatitis C virus, hepatitis C screening, HCV antibody

<https://doi.org/10.21633/jgpha.7.149>

© Merry L. Perry. Originally published in jGPHA (<http://www.gapha.org/jgpha/>) December 20, 2017. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial No-Derivatives License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work ("first published in the Journal of the Georgia Public Health Association...") is properly cited with original URL and bibliographic citation information. The complete bibliographic information, a link to the original publication on <http://www.gapha.jgpha.org/>, as well as this copyright and license information must be included.