

Georgia Southern University

Digital Commons@Georgia Southern

Health and Kinesiology Faculty Publications

Health Sciences and Kinesiology, Department of

6-2017

Parallel and Cross-Sectional Hamstring Injuries in Sprint Running

Li Li

Georgia Southern University, lili@georgiasouthern.edu

Donghai Wang

Shanghai University of Sport

Follow this and additional works at: <https://digitalcommons.georgiasouthern.edu/health-kinesiology-facpubs>



Part of the [Kinesiology Commons](#), and the [Medicine and Health Sciences Commons](#)

Recommended Citation

Li, Li, Donghai Wang. 2017. "Parallel and Cross-Sectional Hamstring Injuries in Sprint Running." *Journal of Sport and Health Science*, 6 (2): 141-142. doi: 10.1016/j.jshs.2017.03.002
<https://digitalcommons.georgiasouthern.edu/health-kinesiology-facpubs/63>

This article is brought to you for free and open access by the Health Sciences and Kinesiology, Department of at Digital Commons@Georgia Southern. It has been accepted for inclusion in Health and Kinesiology Faculty Publications by an authorized administrator of Digital Commons@Georgia Southern. For more information, please contact digitalcommons@georgiasouthern.edu.

Commentary

Parallel and cross-sectional hamstring injuries in sprint running

Li Li ^{a,*}, Donghai Wang ^b

^a School of Health and Kinesiology, Georgia Southern University, Statesboro, GA 30460, USA

^b School of Kinesiology, Shanghai University of Sport, Shanghai 200438, China

Received 21 February 2017; accepted 23 February 2017

Available online 4 March 2017

This issue of *Journal of Sport and Health Science* contains a point-counterpoint discussion of hamstring injuries in sprint running by the groups of Drs. Liu and Yu.^{1–4} They propose different mechanisms of muscle injuries in general, and hamstring injuries in sprint running specifically. Yu et al.^{2,4} present evidence suggesting that hamstring injuries are primarily caused by excessive muscle strain during eccentric contraction. In maximal effort sprint running, excessive muscle strains occur at the end of the swing phase. In contrast, Liu et al.^{1,3} point out that, in addition to excessive hamstring strain, high stresses in the late swing and early stance phase and the transition between swing and stance may also contribute to hamstring injuries when the hamstrings actively assist hip extension and knee flexion.

Both groups provide supporting published, scientific evidence for their contention. Yet, the entire discussion is based on the assumption that the hamstring muscles contract uniformly in all phases of sprint running. However, the different heads of the hamstring muscles have different insertion sites, structure, and fiber type distributions. Therefore, it is safe to assume that the individual heads fulfill different functions and that they do not elongate at the same rate and that stress across them is not uniform.

Using magnetic resonance imaging in combination with finite element modeling, Fiorentino and colleagues⁵ reported that non-uniformity in fiber strains may also be a contributing factor for hamstring injuries, especially when sprinting at high speeds. We propose that this fiber strain and strain rate non-uniformity at fast sprint speeds could lead to “parallel injury”, tissue separation, and misalignment of myofibrils along the muscle fibers (e.g., see Lee and Healy, 2012, for a detailed image⁶). Along these lines, Morgan⁷ proposed the so-called “Popping Sarcomeres Hypothesis”, which is based on the idea that repeated high stresses during eccentric contractions leads

to “tearing” of sarcomeres, and the local damage of sarcomeres leads to more damage in adjacent sarcomeres and neighboring myofibrils due to the sudden increase in localized stress. We propose that the “Popping Sarcomeres” phenomenon could lead to “cross-sectional” hamstring injuries in which the damage occurs tangential to the fiber orientation and produces disruptions of the Z-lines and misaligned A-bands (see Morgan, 1990, for exemplar image⁷). Parallel and cross-sectional hamstring injuries may have different underlying causes and may provide a *post hoc* possibility for evaluating how the injury occurred. In severe hamstring injuries (Grade II or more), one might expect simultaneous parallel and cross-sectional injuries.

The detailed mechanisms of hamstring injuries in sprint running remain a question of debate, and it might be time to look at this issue from an altogether different point of view. Considering that hamstring strains and strain rates might be non-uniform across the different heads, it appears feasible that the non-uniform strains and strain rates may lead to “parallel” injuries, whereas non-uniform stresses might lead to “cross-sectional” injuries. Considering different and multiple hamstring muscle injury mechanisms in sprint running may lead to more targeted training strategies and provide new insights into the prevention and rehabilitation of hamstring injuries.

Authors’ contributions

LL initiated the concept of the paper; LL and DW contributed equally in composition of the essay. Both authors have read and approved the final version of the manuscript, and agree with the order of presentation of the authors.

Competing interests

The authors declare that they have no competing interests.

References

1. Liu Y, Sun Y, Zhu W, Yu J. The late swing and early stance of sprinting are most hazardous for hamstring injuries. *J Sport Health Sci* 2017;6: 133–6.

Peer review under responsibility of Shanghai University of Sport.

* Corresponding author.

E-mail address: lili@georgiasouthern.edu

2. Yu B, Liu H, Garrett WE. Mechanism of hamstring muscle strain injury in sprinting. *J Sport Health Sci* 2017;**6**:130–2.
3. Liu Y, Sun Y, Zhu W, Yu J. Comments to “Mechanism of hamstring muscle strain injury in sprinting” by Yu et al. *J Sport Health Sci* 2017;**6**:139–40.
4. Yu B, Liu H, Garrett WE. Comment on “The late swing and early stance of sprinting are most hazardous for hamstring injuries” by Liu et al. *J Sport Health Sci* 2017;**6**:137–8.
5. Fiorentino NM, Rehorn MR, Chumanov ES, Thelen DG, Blemker SS. Computational models predict larger muscle tissue strains at faster sprinting speeds. *Med Sci Sports Exerc* 2014;**46**:776–86.
6. Lee JC, Healy J. Image of lower limb muscle injury. *Aspetar Sports Med J* 2012;**1**:142–7.
7. Morgan DL. New insights into the behavior of muscle during active lengthening. *Biophys J* 1990;**57**:209–21.