

Georgia Southern University

Georgia Southern Commons

Department of Mathematical Sciences Faculty
Publications

Department of Mathematical Sciences

1-2015

A Family of Multiple Harmonic Sum and Multiple Zeta Star Value Identities

Erin Linebarger
Eckerd College

Jianqiang Zhao
Georgia Southern University, zhaoj8888@gmail.com

Follow this and additional works at: <https://digitalcommons.georgiasouthern.edu/math-sci-facpubs>



Part of the [Education Commons](#), and the [Mathematics Commons](#)

Recommended Citation

Linebarger, Erin, Jianqiang Zhao. 2015. "A Family of Multiple Harmonic Sum and Multiple Zeta Star Value Identities." *Mathematika*, 61 (1): 63-71. doi: 10.1112/S0025579313000302
<https://digitalcommons.georgiasouthern.edu/math-sci-facpubs/413>

This article is brought to you for free and open access by the Department of Mathematical Sciences at Georgia Southern Commons. It has been accepted for inclusion in Department of Mathematical Sciences Faculty Publications by an authorized administrator of Georgia Southern Commons. For more information, please contact digitalcommons@georgiasouthern.edu.

A FAMILY OF MULTIPLE HARMONIC SUM AND MULTIPLE ZETA STAR VALUE IDENTITIES

ERIN LINEBARGER AND JIANQIANG ZHAO

ABSTRACT. In this paper we present a new family of identities for multiple harmonic sums which generalize a recent result of Hessami Pilehrood et al. [7]. We then apply it to obtain a family of identities relating multiple zeta star values to alternating Euler sums. In such a typical identity the entries of the multiple zeta star values consist of blocks of arbitrarily long 2-strings separated by positive integers greater than two while the largest depth of the alternating Euler sums depends only on the number of 2-string blocks but not on their lengths.

1. INTRODUCTION

In a recent paper [7] Hessami Pilehrood et al. proved a few families of identities involving (alternating) multiple harmonic sums and binomial coefficients and, as applications, discovered some new congruences for multiple harmonic sums. In particular, they are able to confirm several conjectures contained in [10] posed by the second author of this paper. They can also provide some new families of identities of MZSV and a new proof of the identity of Zagier [9].

To state one of their results we recall that the (alternating) multiple harmonic sums are multiple variable generalizations of harmonic sums defined by the following: for any $r \in \mathbb{N}$ and $\mathbf{s} = (s_1, s_2, \dots, s_r) \in (\mathbb{Z}^*)^r$

$$H_n(s_1, s_2, \dots, s_r) = \sum_{n \geq k_1 > k_2 > \dots > k_r \geq 1} \prod_{i=1}^r \frac{\operatorname{sgn}(s_i)^{k_i}}{k_i^{|s_i|}}.$$

Throughout the paper we will use \bar{n} to denote a negative entry $s_j = -n$. For example, $H(2, 1) = H(-2, 1)$. Another kind of sums which is closely related to the above is the following star version (also denoted by S in the literature)

$$H_n^*(s_1, s_2, \dots, s_r) = \sum_{n \geq k_1 \geq k_2 \geq \dots \geq k_r \geq 1} \prod_{i=1}^r \frac{\operatorname{sgn}(s_i)^{k_i}}{k_i^{|s_i|}}.$$

We call $l(\mathbf{s}) := r$ the depth of this MHS and $|\mathbf{s}| := \sum_{i=1}^r |s_i|$ the weight. For convenience we set $H_n(\mathbf{s}) = 0$ if $n < l(\mathbf{s})$, $H_n(\emptyset) = H_n^*(\emptyset) = 1$, and $\{s_1, s_2, \dots, s_j\}^d$ the set formed by repeating the composition (s_1, s_2, \dots, s_j) d times. When taking the limit $n \rightarrow \infty$ we get the so-called the (alternating) Euler sum and the (alternating) star Euler sum, respectively:

$$\zeta(\mathbf{s}) = \lim_{n \rightarrow \infty} H_n(\mathbf{s}), \quad \zeta^*(\mathbf{s}) = \lim_{n \rightarrow \infty} H_n^*(\mathbf{s}). \quad (1)$$

When $\mathbf{s} \in \mathbb{N}^\ell$ they are called the *multiple zeta value* (MZV) and the *multiple zeta star value* (MZSV), respectively.

The following is one of the main results of [7].

1991 *Mathematics Subject Classification.* 11A07, 11M32, 11B65, 11B68.

Key words and phrases. Multiple harmonic sum, multiple zeta values, multiple zeta star values, alternating Euler sums.

Theorem 1.1. ([7, Thm. 2.1]) *Let a and b be two non-negative integers. Set $A_{n,k} = (-1)^{k-1} \binom{n}{k} / \binom{n+k}{k}$ for any positive integers n and k . Then for any integer $c \geq 2$,*

$$H_n^*({2}^a, c, {2}^b) = 2 \sum_{k=1}^n \frac{A_{n,k}}{k^{2a+2b+c}} + 4 \sum_{\substack{i+j+|\mathbf{x}|=c \\ i \geq 1, j \geq 2, \mathbf{x} \in \mathbb{N}^r, r \geq 0}} 2^{l(\mathbf{x})} \sum_{k=1}^n \frac{H_{k-1}(\mathbf{x}, i+2b) A_{n,k}}{k^{2a+j}},$$

where in the sum $i \geq 1, j \geq 2$ and \mathbf{x} runs through all possible compositions of positive integers (or the empty set when $r = 0$) satisfying the restriction $i + j + |\mathbf{x}| = c$.

In sections 2 and 3 we generalize Theorem 1.1 to express $H_n^*(\mathbf{s})$ using binomial coefficients where $\mathbf{s} = (\{2\}^{a_1}, c_1, \dots, \{2\}^{a_r}, c_r, \{2\}^{a_{r+1}})$ with positive integers $c_1, \dots, c_r \geq 2$. Taking limit and using a key lemma we shall obtain a new family of identities of MZSV involving alternating Euler sums in the following Theorem (an equivalent form of Theorem 4.1).

Theorem 1.2. *Let n be a positive integer and $\mathbf{s} = (\{2\}^{a_1}, c_1, \dots, \{2\}^{a_r}, c_r, \{2\}^{a_{r+1}})$ where $a_j, c_j \in \mathbb{N}_0$ and $c_j \geq 3$. Then we have*

$$\zeta^*(\mathbf{s}) = - \sum_{\mathbf{p}=(\overline{2a_1+2}) \circ 1^{\circ(c_1-3)} \circ (2a_2+3) \circ 1^{\circ(c_2-3)} \circ \dots \circ (2a_r+3) \circ 1^{\circ(c_r-3)} \circ (2a_{r+1}+1)}$$

where \circ is either the comma “,” or the O-plus “ \oplus ” defined by $\alpha \oplus \beta = \text{sgn}(\alpha) \text{sgn}(\beta) (|\alpha| + |\beta|)$ for all $\alpha, \beta \in \mathbb{Z}^*$, and $1^{\circ c} = \underbrace{1 \circ \dots \circ 1}_c$.

For example,

$$\begin{aligned} \zeta^*({2}^7, 3, {2}^2, 3, {2}^3) &= -8\zeta(\overline{16}, 7, 7) - 4\zeta(\overline{23}, 7) - 4\zeta(\overline{16}, 14) - 2\zeta(\overline{30}), \\ \zeta^*({2}^3, 3, {2}^2, 4, {2}^5) &= -16\zeta(\overline{8}, 7, 1, 11) - 8\zeta(\overline{8}, 7, 12) - 8\zeta(\overline{8}, 8, 11) \\ &\quad - 8\zeta(\overline{15}, 1, 11) - 4\zeta(\overline{15}, 12) - 4\zeta(\overline{16}, 11) - 4\zeta(\overline{8}, 19) - 2\zeta(\overline{27}). \end{aligned}$$

Broadhurst considered alternating Euler sums as “honorary” MZVs and relates them to the study of knots and Feynman diagrams (see [1]). From the point of view of algebraic geometry, Goncharov and Manin [5] showed that the MZVs and MZSVs are periods of mixed Tate motives over \mathbb{Z} (see also [2]) while Deligne and Goncharov [4] proved that the alternating Euler sums are periods of mixed Tate motives over \mathbb{Z} ramified at 2 (see also [3]). This provides a kind of inverse to “Galois descent” since the linear combinations of the Euler sums in the above examples are invariant under a certain motivic Galois group. In general it is not easy to show this unramifiedness property directly. Another important aspect of Theorem 1.2 is that the depth of the MZSV can be very large when the 2-strings are very long, but the largest depth of the Euler sums on the right hand side depends only on the number of such strings but not on their lengths.

The ideas used in this paper have been applied to obtain similar results for other types of strings in [11]. In particular, the conjectural Two-one formula of Ohno and Zudilin is proved there. Its q -analog has been obtained by Hessami Pilehroods recently in [6]. In [8] Hessami Pilehroods and the second author further prove a q -analog of the main result of this paper (Theorem 2.3) and some similar results for other types of strings contained in [11].

2. IDENTITIES FOR MULTIPLE HARMONIC SUMS

The following lemma is the special case of [7, Lemma 2.2] when $m = 2$ and $c_n^{(2)} = n!^2/(2n)!$.

Lemma 2.1. For any positive integers n and k define $A_{n,k} = (-1)^{k-1} \binom{n}{k} / \binom{n+k}{k}$. Then for every $c \in \mathbb{N}$ and $\mathbf{v} \in \mathbb{N}^t$ ($t \geq 0$) we have

$$\frac{1}{n^c} \sum_{k=1}^n \frac{H_{k-1}(\mathbf{v}) A_{n,k}}{k^a} = \sum_{k=1}^n \frac{H_{k-1}(\mathbf{v}) A_{n,k}}{k^{a+c}} + \sum_{\substack{j+|\mathbf{x}|=a+c \\ j \geq 0, x_r > a}} 2^{l(\mathbf{x})} \sum_{k=1}^n \frac{H_{k-1}(\mathbf{x}, \mathbf{v}) A_{n,k}}{k^j}$$

where $\mathbf{x} = (x_1, x_2, \dots, x_r) \in \mathbb{N}^r$ for $r \geq 0$, and $|\mathbf{x}| = \sum_{i=1}^r x_i$, $l(\mathbf{x}) = r$. Here $|\mathbf{x}| = 0$ if $r = 0$ which implies that $\mathbf{x} = \emptyset$.

Definition 2.2. Suppose $\mathbf{s} = (\{2\}^{a_1}, c_1, \dots, \{2\}^{a_r}, c_r, \{2\}^{a_{r+1}})$. We define two kinds of string operations on the condensation $\tilde{\mathbf{s}} := (2a_1, c_1, \dots, 2a_r, c_r, 2a_{r+1})$ of \mathbf{s} as follows.

- A merge μ_t for some $1 \leq t \leq r$ changes the symbols “ c_t ,” to “ $+c_t+$ ”,
- A substitution σ_t changes the symbols “ c_t ,” to “ $+j_t, \mathbf{x}_t, i_t+$ ”.

Let $I = \{i_1, \dots, i_t\}$ be a subset of $[r] := \{1, \dots, r\}$. Define

- The merge operation $\mu_I = \mu_{i_1} \circ \dots \circ \mu_{i_t}$ and similarly for substitutions σ_I ,
- The complement $\bar{I} = [r] \setminus I$,
- The combined operation $\kappa_I = \sigma_{\bar{I}} \circ \mu_I$.

For any composition (e_1, \dots, e_r) of natural numbers we denote

- The first component $e_1 = \varphi(\mathbf{e})$
- The truncation operator $\text{Tr}(e_1, \dots, e_r) = (e_2, \dots, e_r)$.
- The negation operator $\nu(e_1, \dots, e_r) = (\bar{e}_1, e_2, \dots, e_r)$.

Notice that if $r = 1$ then $\text{Tr}(e_1) = \emptyset$.

Theorem 2.3. Let n be a positive integer and $\mathbf{s} = (\{2\}^{a_1}, c_1, \dots, \{2\}^{a_r}, c_r, \{2\}^{a_{r+1}})$ where $a_j, c_j \in \mathbb{N}_0$ and $c_j \geq 2$. Set $A_{n,k} = (-1)^{k-1} \binom{n}{k} / \binom{n+k}{k}$. Then

$$H_n^*(\mathbf{s}) = 2 \sum_{I \subseteq [r]} \sum_{\substack{i_t + j_t + |\mathbf{x}_t| = c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^n \frac{H_{k-1}(\text{Tr} \circ \kappa_I(\tilde{\mathbf{s}})) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{s}})}}, \quad (2)$$

where $\tilde{\mathbf{s}} = (2a_1, c_1, \dots, 2a_r, c_r, 2a_{r+1})$ is the condensation of \mathbf{s} .

Example 2.4. When $r = 1$ we have two possible subsets of $[1]$: $I = \emptyset$ and $I = [1]$. our Theorem 2.3 becomes Theorem 1.1. When $r = 2$ we get the following: For a positive integer n and non-negative integers a_1, a_2, a_3 and positive integers $c_1, c_2 \geq 2$ we set $\mathbf{s} = (\{2\}^{a_1}, c_1, (\{2\}^{a_2}, c_2, (\{2\}^{a_3}))$. Then

$$\begin{aligned} \kappa_{[2]} &= (2a_1 + c_1 + 2a_2 + c_2 + 2a_3), & \kappa_{\emptyset} &= (2a_1 + j_1, \mathbf{x}_1, i_1 + 2a_2 + j_2, \mathbf{x}_2, i_2 + 2a_3), \\ \kappa_{\{1\}} &= (2a_1 + c_1 + 2a_2 + j_2, \mathbf{x}_2, i_2 + 2a_3), & \kappa_{\{2\}} &= (2a_1 + j_1, \mathbf{x}_1, i_1, 2a_2 + c_2 + 2a_3). \end{aligned}$$

Hence by Theorem 2.3

$$\begin{aligned}
H_n^*(\mathbf{s}) &= 2 \sum_{k=1}^n \frac{A_{n,k}}{k^{2a_1+c_1+2a_2+c_2+2a_3}} + 4 \sum_{\substack{i+j+|\mathbf{x}|=c_2 \\ i \geq 1, j \geq 2}} 2^{l(\mathbf{x})} \sum_{k=1}^n \frac{H_{k-1}(\mathbf{x}, i+2a_3) A_{n,k}}{k^{2a_1+c_1+2a_2+j}} \\
&+ 8 \sum_{\substack{i_1+j_1+|\mathbf{x}_1|=c_1, i_2+j_2+|\mathbf{x}_2|=c_2 \\ i_1, i_2 \geq 1, j_1, j_2 \geq 2}} 2^{l(\mathbf{x}_1)+l(\mathbf{x}_2)} \sum_{k=1}^n \frac{H_{k-1}(\mathbf{x}_1, i_1+2a_2+j_2, \mathbf{x}_2, i_2+2a_3) A_{n,k}}{k^{2a_1+j_1}} \\
&+ 4 \sum_{\substack{i+j+|\mathbf{x}|=c_1 \\ i \geq 1, j \geq 2}} 2^{l(\mathbf{x})} \sum_{k=1}^n \frac{H_{k-1}(\mathbf{x}, i+2a_2+c_2+2a_3) A_{n,k}}{k^{2a_1+j}}.
\end{aligned}$$

3. PROOF OF THEOREM 2.3

We proceed by induction on the sum $r+n$. Since $r \geq 0, n \geq 1$, our base case is $r+n=1$ which implies $H_1^*(\{2\}^a) = 1$, and therefore the formula is true. Suppose the statement is true when $r+n=k-1$, and let $r+n=k$. Then by definition

$$H_n^*(\mathbf{s}) = \sum_{l=0}^{a_1} \frac{1}{n^{2(a_1-l)}} H_{n-1}^*(\mathbf{t}_l) + \frac{1}{n^{2a_1+c_1}} H_n^*(\mathbf{u}),$$

where $\mathbf{t}_l = (\{2\}^l, c_1, \{2\}^{a_2}, c_2, \dots, c_r, \{2\}^{a_{r+1}})$ and $\mathbf{u} = (\{2\}^{a_2}, c_2, \dots, c_r, \{2\}^{a_{r+1}})$. We now set their condensations as follows:

$$\tilde{\mathbf{t}}_l = (2l, c_1, 2a_2, c_2, \dots, c_r, 2a_{r+1}), \quad \tilde{\mathbf{u}} = (2a_2, c_2, \dots, c_r, 2a_{r+1}).$$

By inductive hypothesis,

$$\begin{aligned}
H_n^*(\mathbf{s}) &= \sum_{l=0}^{a_1} \frac{2}{n^{2(a_1-l)}} \sum_{I \subseteq [r]} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}|+\sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^{n-1} \frac{H_{k-1}(\text{Tr} \circ \kappa_I(\tilde{\mathbf{t}}_l)) A_{n-1,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{t}}_l)}} \\
&+ \frac{2}{n^{2a_1+c_1}} \sum_{I \subseteq [r] \setminus \{1\}} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}|+\sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^n \frac{H_{k-1}(\text{Tr} \circ \kappa_I(\tilde{\mathbf{u}})) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{u}})}}.
\end{aligned}$$

Changing the order of summation and summing the inner sum we get

$$A_{n-1,k} \sum_{l=0}^{a_1} \frac{n^{2l}}{k^{2l}} = (-1)^{k-1} \frac{n^{2a_1+2} - k^{2a_1+2}}{(n^2 - k^2)k^{2a_1}} \frac{\binom{n-1}{k}}{\binom{n-1+k}{k}} = \left(\frac{n^{2a_1}}{k^{2a_1}} - \frac{k^2}{n^2} \right) A_{n,k}.$$

Thus

$$\begin{aligned}
H_n^*(\mathbf{s}) &= 2 \sum_{I \subseteq [r]} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}|+\sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^n \frac{H_{k-1}(\text{Tr} \circ \kappa_I(\tilde{\mathbf{t}}_l)) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{t}}_l)-2l+2a_1}} \\
&- \frac{2}{n^{2a_1+2}} \sum_{I \subseteq [r]} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}|+\sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^n \frac{H_{k-1}(\text{Tr} \circ \kappa_I(\tilde{\mathbf{t}}_l)) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{t}}_l)-2l-2}}
\end{aligned}$$

$$+ \frac{2}{n^{2a_1+c_1}} \sum_{I \subseteq [r] \setminus \{1\}} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^n \frac{H_{k-1} \left(\text{Tr} \circ \kappa_I(\tilde{\mathbf{u}}) \right) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{u}})}}.$$

Since $\tilde{\mathbf{t}}_l$ coincides with $\tilde{\mathbf{u}}$ everywhere except for the first component, $\text{Tr} \circ \kappa_I(\tilde{\mathbf{t}}_l) = \text{Tr} \circ \kappa_I(\tilde{\mathbf{u}})$. Similarly, $\varphi \circ \kappa_I(\tilde{\mathbf{t}}_l) - 2l + 2a_1 = \varphi \circ \kappa_I(\tilde{\mathbf{u}})$. Hence, by induction we only need to show the following quantity vanishes:

$$\begin{aligned} & \frac{2}{n^{2a_1+c_1}} \sum_{I \subseteq [r] \setminus \{1\}} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^n \frac{H_{k-1} \left(\text{Tr} \circ \kappa_I(\tilde{\mathbf{u}}) \right) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{u}})}} \\ & - \frac{2}{n^{2a_1+2}} \sum_{I \subseteq [r]} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^n \frac{H_{k-1} \left(\text{Tr} \circ \kappa_I(\tilde{\mathbf{t}}_l) \right) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{t}}_l) - 2l - 2}} \\ & = \frac{2}{n^{2a_1+c_1}} \sum_{I \subseteq [r] \setminus \{1\}} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^n \frac{H_{k-1} \left(\text{Tr} \circ \kappa_I(\tilde{\mathbf{u}}) \right) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{u}})}} \\ & - \frac{2}{n^{2a_1+2}} \sum_{I \subseteq [r] \setminus \{1\}} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^n \frac{H_{k-1} \left(\text{Tr} \circ \kappa_I(\tilde{\mathbf{t}}_l) \right) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{t}}_l) - 2l - 2}} \\ & - \frac{4}{n^{2a_1+2}} \sum_{I \subseteq [r] \setminus \{1\}} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^n \frac{H_{k-1} \left(\text{Tr} \circ \kappa_{I \cup \{1\}}(\tilde{\mathbf{t}}_l) \right) A_{n,k}}{k^{\varphi \circ \mu_{\bar{I}} \circ \sigma_{I \cup \{1\}}(\tilde{\mathbf{t}}_l) - 2l - 2}} \\ & = \frac{2}{n^{2a_1+2}} \sum_{I \subseteq [r] \setminus \{1\}} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \left(\frac{1}{n^{c_1-2}} \sum_{k=1}^n \frac{H_{k-1} \left(\text{Tr} \circ \kappa_I(\tilde{\mathbf{u}}) \right) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{u}})}} \right. \\ & \left. - 2 \sum_{\substack{i_1+j_1+|\mathbf{x}_1|=c_1, \\ i_1 \geq 1, j_1 \geq 2}} 2^{l(\mathbf{x}_1)} \sum_{k=1}^n \frac{H_{k-1} \left(\text{Tr} \circ \kappa_I(\tilde{\mathbf{t}}_l) \right) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{t}}_l) - 2l - 2}} - \sum_{k=1}^n \frac{H_{k-1} \left(\text{Tr} \circ \kappa_{I \cup \{1\}}(\tilde{\mathbf{t}}_l) \right) A_{n,k}}{k^{\varphi \circ \kappa_{I \cup \{1\}}(\tilde{\mathbf{t}}_l) - 2l - 2}} \right). \quad (3) \end{aligned}$$

Note that if $1 \notin I$ then $\text{Tr} \circ \kappa_I(\tilde{\mathbf{u}}) = \text{Tr} \circ \kappa_{I \cup \{1\}}(\tilde{\mathbf{t}}_l)$, which we will now denote by \mathbf{v} . Furthermore, $\varphi \circ \kappa_{I \cup \{1\}}(\tilde{\mathbf{t}}_l) = \varphi \circ \kappa_I(\tilde{\mathbf{u}}) + c_1 + 2l$. Let $a = \varphi \circ \kappa_I(\tilde{\mathbf{u}})$. Then

$$\kappa_I(\tilde{\mathbf{t}}_l) = \sigma_{\bar{I}} \circ \mu_I(\tilde{\mathbf{t}}_l) = \sigma_1(2l, c_1, a, \dots) = (j_1 + 2l, \mathbf{x}_1, i_1 + a, \dots).$$

Set $\mathbf{w} = (\mathbf{x}_1, i_1 + a)$. Then we see that the three sums inside the parenthesis (3) can be rewritten as

$$\frac{1}{n^{c_1-2}} \sum_{k=1}^n \frac{H_{k-1}(\mathbf{v}) A_{n,k}}{k^a} - \sum_{\substack{j_1+|\mathbf{w}|=c_1+a_1, \\ j_1 \geq 2, w_p > a_1}} 2^{l(\mathbf{w})} \sum_{k=1}^n \frac{H_{k-1}(\mathbf{w}, \mathbf{v}) A_{n,k}}{k^{j_1-2}} - \sum_{k=1}^n \frac{H_{k-1}(\mathbf{v}) A_{n,k}}{k^{a+c_1-2}},$$

where $w_p = i + a_1$ ($i \geq 1$) is the last component of \mathbf{w} . This last expression vanishes by taking $c = c_1 - 2$ and $j = j_1 - 2$ in Lemma 2.1. We have now completed the proof of Theorem 2.3.

4. THE KEY LEMMA AND AN IDENTITY FAMILY OF MZSV

In [7], using the corresponding identities of MHS Hessami Pilehrood et al. find some new families of identities of MZSV and a new proof of the identity of Zagier [9]. We can similarly use Theorem 2.3 to obtain a new family of MZSV as follows.

Theorem 4.1. *Let n be a positive integer and $\mathbf{s} = (\{2\}^{a_1}, c_1, \dots, \{2\}^{a_r}, c_r, \{2\}^{a_{r+1}})$ where $a_j, c_j \in \mathbb{N}_0$ and $c_j \geq 2$. Set $\tilde{\mathbf{s}} = (2a_1, c_1, \dots, 2a_r, c_r, 2a_{r+1})$. Then with notation given by Definition 2.2 we have*

$$\zeta^*(\mathbf{s}) = 2 \sum_{I \subseteq [r]} \sum_{\substack{i_t + j_t + |\mathbf{x}_t| = c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \zeta\left(\nu \circ \kappa_I(\tilde{\mathbf{s}})\right). \quad (4)$$

Proof. By taking limit $n \rightarrow \infty$ in Theorem 2.3 we see that (4) follows immediately from the following lemma. \square

Lemma 4.2. *Let $d \in \mathbb{N}_0$ and let e be a real number with $e > 1$. Then for all $\mathbf{s} \in (\mathbb{Z}^*)^d$ ($\mathbf{s} = \emptyset$ if $d = 0$) we have*

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{|H_{k-1}(\mathbf{s})|}{k^e} \left(1 - \frac{\binom{n}{k}}{\binom{n+k}{k}}\right) = 0. \quad (5)$$

Remark 4.3. This is the key step which enables us to go from MHS identities to MZSV identities in this paper. Apparently the authors of [7] have already used this result in their paper although no proof is given there. Because of its importance we provide the following detailed analysis. Notice that for any fixed k we have $\lim_{n \rightarrow \infty} \binom{n}{k} / \binom{n+k}{k} = 1$ but for k close to n , say $k = n$, we have $\lim_{n \rightarrow \infty} \binom{n}{k} / \binom{n+k}{k} = 0$ by Stirling's formula. So (5) is not obvious to us.

Proof. First we have

$$\begin{aligned} 0 < 1 - \frac{\binom{n}{k}}{\binom{n+k}{k}} &= 1 - \frac{n(n-1)(n-2)\cdots(n-k+1)}{(n+1)(n+2)\cdots(n+k)} \\ &= 1 - \frac{n(n-1)\left(\frac{n}{2}-1\right)\cdots\left(\frac{n}{k-1}-1\right)}{k(n+1)\left(\frac{n}{2}+1\right)\cdots\left(\frac{n}{k}+1\right)} \\ &= 1 - \frac{n^k H_{k-1}(\{1\}^{k-1}) - n^{k-1} H_{k-1}(\{1\}^{k-2}) + \cdots + (-1)^k n^2 H_{k-1}(1) - n(-1)^k}{k(n^k H_k(\{1\}^k) + n^{k-1} H_k(\{1\}^{k-1}) + \cdots + n H_k(1) + 1)} \end{aligned} \quad (6)$$

$$\begin{aligned} &\leq \frac{n^{k-1} H_k(\{1\}^{k-2}) + n^{k-2} H_k(\{1\}^{k-3}) + \cdots + n^2 H_k(1) + n}{k(n^k H_k(\{1\}^k) + n^{k-1} H_k(\{1\}^{k-1}) + \cdots + n H_k(1) + 1)} \\ &\quad + \frac{n^{k-1} H_k(\{1\}^{k-1}) + \cdots + n H_k(1) + 1}{n^k H_k(\{1\}^k) + n^{k-1} H_k(\{1\}^{k-1}) + \cdots + n H_k(1) + 1}. \end{aligned} \quad (7)$$

Here from (6) to (7) we canceled the leading term of the numerator in (6) because of $H_k(\{1\}^k) = 1/k!$, took absolute values of each term on the right and then changed all

H_{k-1} to H_k . Observe that for any $1 \leq j \leq k$ we have

$$\begin{aligned} H_k(\{1\}^j) &= \sum_{k \geq n_1 > \dots > n_j > 0} \frac{1}{n_1 \cdots n_j} \\ &\leq \sum_{k \geq n_1 > \dots > n_j > n_{j+1} > 0} \frac{k}{n_1 \cdots n_j \cdot n_{j+1}} = k H_k(\{1\}^{j+1}) \leq \dots \leq k^{k-j} H_k(\{1\}^k), \end{aligned}$$

which is even true for $j = 0$. Thus we get

$$1 - \frac{\binom{n}{k}}{\binom{n+k}{k}} \leq \frac{2(n^{k-1}k + n^{k-2}k^2 + \dots + nk^{k-1} + k^k)}{n^k} \leq \frac{2k^2}{n}.$$

Now for any $\mathbf{s} \in (\mathbb{Z}^*)^d$ we clearly have

$$|H_{k-1}(\mathbf{s})| \leq H_{k-1}(\{1\}^d) \leq C \log^d(k)$$

for some positive constant C . Therefore by setting $\delta = \min\{(e-1)/2, 1/2\} > 0$ we have

$$\sum_{k=1}^n \frac{|H_{k-1}(\mathbf{s})|}{k^e} \left(1 - \frac{\binom{n}{k}}{\binom{n+k}{k}}\right) \leq \frac{2C}{n} \sum_{k=1}^{\lfloor n^\delta \rfloor} \frac{\log^d(k)}{k^{e-2}} + C \sum_{k=\lfloor n^\delta \rfloor + 1}^{\infty} \frac{\log^d(k)}{k^e}. \quad (8)$$

If $e \geq 2$ then

$$\frac{2C}{n} \sum_{k=1}^{\lfloor n^\delta \rfloor} \frac{\log^d(k)}{k^{e-2}} \leq \frac{2C}{n} \sum_{k=1}^{\sqrt{n}} \log^d(k) \leq \frac{2C \log^d(\sqrt{n})}{\sqrt{n}} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

If $1 < e < 2$ then

$$\frac{2C}{n} \sum_{k=1}^{\lfloor n^\delta \rfloor} \frac{\log^d(k)}{k^{e-2}} \leq \frac{2C n^{\delta(3-e)} \log^d(n^\delta)}{n} \leq \frac{2C \delta^d \log^d(n)}{\sqrt{n}} \rightarrow 0 \text{ as } n \rightarrow \infty,$$

since $1 + \delta(e-3) = 1 + (e-1)(e-3)/2 = (e-2)^2/2 + 1/2 \geq 1/2$. Now the last sum on the right of (8) is the tail of a convergent series so (5) follows immediately and therefore the lemma is proved. \square

Acknowledgement. We would like to thank Roberto Tauraso for sending us their preprint [7] upon which we can build our current work, Wadim Zudilin for kindly showing us an improvement on Lemma 4.2, and K. Hessami Pilehrood, T. Hessami Pilehrood and Y. Ohno for pointing out some misprints in the first version of this paper. We thank the anonymous referees who provided a few very insightful remarks which are incorporated in the paper. JZ is supported by an NSF grant DMS1162116.

REFERENCES

- [1] D.J. Broadhurst, Conjectured enumeration of irreducible multiple zeta values, from knots and Feynman diagrams, preprint, [arXiv:hep-th/9612012](https://arxiv.org/abs/hep-th/9612012)
- [2] F. Brown, Mixed Tate motives over $\text{Spec}(\mathbb{Z})$, *Ann. Math.* **175**(2) (2012), pp. 949–976.
- [3] P. Deligne, Le groupe fondamental de la $\mathbb{G}_m - \mu_N$, pour $N = 2, 3, 4, 6$ ou 8 , *Publ. Math. Inst. Hautes Etudes Sci.* **112** (2010), pp. 101–141.
- [4] P. Deligne and A. Goncharov, Groupes fondamentaux motiviques de Tate mixte, *Ann. Sci. Ecole Norm. S.* **38**(1) (2005), pp. 1–56.
- [5] A.B. Goncharov and Y.I. Manin, Multiple ζ -motives and moduli spaces $M_{0,n}$, *Compositio Math.* **140** (2004), pp. 1–14.
- [6] Kh. Hessami Pilehrood and T. Hessami Pilehrood, On q -analogues of two-one formulas for multiple harmonic sums and multiple zeta star values, preprint [arXiv: 1304.0269\[NT\]](https://arxiv.org/abs/1304.0269).

- [7] Kh. Hessami Pilehrood, T. Hessami Pilehrood and R. Tauraso, New properties of multiple harmonic sums modulo p and p -analogues of Leshchiner's series, to appear in *Trans. Amer. Math. Soc.*, DOI: <http://dx.doi.org/10.1090/S0002-9947-2013-05980-6>, arXiv:1206.0407
- [8] Kh. Hessami Pilehrood, T. Hessami Pilehrood and J. Zhao, On q -analogues of some families of multiple harmonic sum and multiple zeta star value identities, arXiv: 1307.7985
- [9] D. Zagier, Evaluation of the multiple zeta values $\zeta(2, \dots, 2, 3, 2, \dots, 2)$, *Ann. of Math.* **175** (2012), pp. 977–1000.
- [10] J. Zhao, Wolstenholme type Theorem for multiple harmonic sums, *Int. J. of Number Theory* **4** (2008), no. 1, pp. 73–106, arXiv:math.NT/0301252
- [11] J. Zhao, Identity families of multiple harmonic sums and multiple zeta (star) values, preprint arXiv:1303.2227

DEPARTMENT OF MATHEMATICS, ECKERD COLLEGE, ST. PETERSBURG, FL 33711

KAVLI INSTITUTE FOR THEORETICAL PHYSICS CHINA, BEIJING, CHINA, MAX PLANCK INSTITUTE FOR MATHEMATICS, VIVATSGASSE 7, 53111 BONN, GERMANY, AND DEPARTMENT OF MATHEMATICS, ECKERD COLLEGE, ST. PETERSBURG, FL 33711

E-mail address: emlineba@eckerd.edu

E-mail address: zhaoj@eckerd.edu