

Fall 2007

Phylogenetic Relationships within Coleeae (Bignoniaceae Juss.)

Andreina Carvajal

Follow this and additional works at: <https://digitalcommons.georgiasouthern.edu/etd>

Recommended Citation

Carvajal, Andreina, "Phylogenetic Relationships within Coleeae (Bignoniaceae Juss.)" (2007). *Electronic Theses and Dissertations*. 697.
<https://digitalcommons.georgiasouthern.edu/etd/697>

This thesis (open access) is brought to you for free and open access by the Jack N. Averitt College of Graduate Studies at Georgia Southern Commons. It has been accepted for inclusion in Electronic Theses and Dissertations by an authorized administrator of Georgia Southern Commons. For more information, please contact digitalcommons@georgiasouthern.edu.

PHYLOGENETIC RELATIONSHIPS WITHIN COLEEAE (BIGNONIACEAE JUSS.)

by

ANDREINA FUENTES CARVAJAL

(Under the Direction of Michelle Zjhra)

ABSTRACT

On Madagascar, Bignoniaceae is represented by tribes Coleeae (4 genera/58 species) and Tecomeae (4 genera/15 species). Species of Coleeae occur in assemblages of sympatric, locally endemic species. The purpose of this study was to explore the phylogenetic relationships within Coleeae. In order to do so, techniques such as PCR by using universal primers for *waxy* nuclear gene, agarose electrophoresis, cloning and sequencing were performed. A total of 26 taxa were successfully amplified. Sequences were analyzed using three different phylogenetic programs (Maximum parsimony, maximum likelihood and Bayesian analysis) resulting in various phylogenetic trees. Main conclusions drawn from this work are: 1. *waxy* is a good molecular marker to resolve relationships within Coleeae; 2. ITS and *waxy* resolved main clades similarly showing monophyly of *Colea* and *Rhodocolea*. Since conservation requires reliable information on species identity, a good phylogenetic assessment is of great importance in determining taxonomic status of organisms.

INDEX WORDS: Bignoniaceae, Coleeae, Phylogeny, GBSSI, *waxy*.

PHYLOGENETIC RELATIONSHIP WITHIN COLEEEAE (BIGNONIACEAE JUSS.)

by

ANDREINA FUENTES CARVAJAL

B. S., Universidad de Oriente, Venezuela, 2003

A Thesis Submitted to the Graduate Faculty of Georgia Southern University in Partial
Fulfillment of the Requirements for the Degree

MASTER OF SCIENCE

STATESBORO, GA

2007

© 2007

Andreina Fuentes Carvajal

All Rights Reserved

PHYLOGENETIC RELATIONSHIP WITHIN COLEEEAE (BIGNONIACEAE JUSS.)

by

ANDREINA FUENTES CARVAJAL

Major Professor:	Michelle Zjhra
Committee:	Lorenza Beati
	James Hutcheon

Electronic Version Approved:

December 2007

DEDICATION

To my mother, Morella, my inspiration for everything I do.

To Michelle Zjhra and Tatjana Good, for believing in me.

To my husband, Matthew, because of the happiness you have brought into my life.

ACKNOWLEDGMENTS

To my advisor, Dr. Michelle Zjhra

To my committee, Dr. James Hutcheon and Dr. Lorenza Beati,

To the Graduate students office,

To the Irene Boole foundation,

To Dr. Laura Regassa,

To Dr. Chris Cuttler,

To my friends Tatjana Good and Pete Way.

To Dr. Ray Chandler,

To Dr. Steve Vives, and

To Dr. Dick Diebolt

TABLE OF CONTENTS

	Page
ACKNOWLEDGMENTS.....	6
LIST OF TABLES.....	8
LIST OF FIGURES.....	9
CHAPTER	
1 INTRODUCTION.....	10
2 MATERIAL AND METHODS.....	14
3 RESULTS.....	21
4 DISCUSSION.....	34
5 LITERATURE CITED.....	38

LIST OF TABLES

	Page
Table 1: PCR's general conditions.....	19
Table 2: Primers used in PCR amplification and sequencing of <i>waxy</i>	19
Table 3: Summarized results of modeltest.....	33

LIST OF FIGURES

	Page
Figure 1: Diagram of granule-bound synthase gene from <i>Solanum tuberosum</i>	20
Figure 2. Map of distribution of Coleeae's species	24
Figure 3. Phylogenetic trees from analyses of ITS	27
Figure 4. Phylogenetic trees from analysis of a subset of <i>waxy</i>	28
Figure 5. Phylogenetic trees from analysis of <i>waxy</i>	29
Figure 6. Phylogenetic trees from analysis of combined data 1 (ITS+ <i>waxy</i>).	30
Figure 7. Phylogenetic trees from analysis of combined data 2 (ITS+subset <i>waxy</i>)...	31
Figure 8. Bayesian phylogenetic trees	32

CHAPTER 1

INTRODUCTION

Taxonomy of Coleeae.

Bignoniaceae (the Trumpet Creeper family) is predominately a tropical family of trees and lianas. The family is characterized by opposite, compound leaves; flowers that are five-merous, fused, and zygomorphic with four stamens; fruit that are usually elongate, septicidal to loculicidal capsules (De Jussie, 1789). Phylogenetic studies of the family have placed Bignoniaceae in order Lamiales, within Euasterids I of the core Eudicots (Olmstead *et al.*, 1993, APG II, 2003).

The island of Madagascar is home to the world's second greatest center of diversity for Bignoniaceae, exceeded only by the much larger neotropical region (Gentry, 1988). The Malagasy diversity of Bignoniaceae is represented by near endemic tribe Coleeae (4 genera/58 species) and pantropical Tecomeae (4 genera/15 species) (Zjhra, 2006).

The first classification to consider Madagascar species was that of Bojer (1837) who divided the family into three tribes: Bignonieae, Crescentieae and Coleeae. Further classifications considered Coleeae as part of Crescentieae (De Candolle, 1838; 1845; Seeman 1860; Baillon 1887, 1888; Perrier De la Bathie, 1938) due predominately to the shared but unusual characteristics of indehiscent, fleshy, animal-dispersed fruit. Gentry (1976, 1980) circumscribed Bignoniaceae into 7 tribes: Coleeae, Bignonieae, Crescentieae, Tecomeae, Eccremocarpeae, Tourrettiae, and Oroxyleae. Furthermore, Gentry (1980) suggested that similarities between Coleeae and Crescentieae were due to derived position of Bignoniaceae among the angiosperms means that Bignoniaceae

arrived on Madagascar via long-distance dispersal. The late arrival of Coleeae, compared to earlier Angiosperm groups, suggest that species of Coleeae radiated rapidly and recently on Madagascar (Gentry, 1980). Coleeae, therefore, make an ideal group for studies of diversification.

Species of Coleeae occur in sympatric assemblages and are often local endemics (Zjhra, 1998, Good *et al.*, 2006, Zjhra, in review). Zjhra (2003) investigated the radiation and diversification of traits using a nrDNA (ITS) derived phylogeny and found that traditional generic delimitations were not monophyletic. Yet cpDNA (ndhF, trnT-L, trnL-F spacer data) derived relationships suggested monophyletic generic clades (Zjhra *et al.*, 2004). The latter could be possible due to the slower rates of evolution of chloroplast genome in comparisons to nuclear DNA, which have make them useful in resolving higher taxonomic levels than species (Sang, 2002).

Molecular markers used to resolve phylogenies

Plant molecular systematists predominately have relied upon chloroplast DNA (cpDNA) and nuclear highly repetitive rDNA arrays, using conserved genes to study ancient divergence events and spacer regions for lower taxonomic level (Mason-Gamer *et al.* 1998). However, cpDNA is generally inherited as a single unit and lacks recombination, thus phylogenies based on cpDNA effectively represent a single gene tree (Doyle, 1992) that traces the genealogy of one parent (Sang, 2002).

The internal transcribed spacer (ITS) is the most frequently sequenced nuclear region due to its comparatively easy amplification, compared to other nDNA markers, and sufficient variability to resolve phylogenetic relationship in different plants groups at the generic level or lower (Mort & Crawford, 2004). However, poor resolution at lower

taxonomic levels (Baldwin *et al.*, 1995) is likely due to lack of sufficient informative characters in the short length of the spacer (Sang, 2002) in some groups.

The need for additional molecular markers for reconstructing speciation events is fundamentally a need for constructing organismal phylogenies rather than single gene trees (Cronn *et al.*, 2003, Doyle, 1992). Low-copy genes, such as GBSSI, have attracted the attention of systematists since they offer several advantages such as limited concerted evolution (Álvarez & Wendel, 2003). Problems, however, about the use of low-copy loci include marker selection, primer design, duplications, and deletions, which can potentially lead to the reconstruction of gene duplication events, rather than speciation events (Sang, 2002).

The granule-bound starch synthase (GBSSI) gene, also known as *waxy* (Echt & Schwartz, 1981), encodes an enzyme involved in starch synthesis. This starch synthesis enzyme adds glucose residue from ADP glucose to the non-reducing end of a growing glucan chain and is the only one within the starch synthases that is required for amylose synthesis, although there is some evidence of glucose incorporation into amylopectin in starch granules (Denyer *et al.*, 2001). The *waxy* gene cloned in potato (van der Leij *et al.* 1991) comprised 4663 bp; the translated portion encompassed 2961, had 13 translated exons, and 13 introns, without the one in the leader. It encoded a 58.2 kilodalton mature protein with 540 amino acids. *Waxy* is a low-copy nuclear gene, existing as a single copy in nearly all plants examined (Mason & Gamer *et al.*, 1998). The exceptions to date, however, are duplications in Rosaceae and Rhamnaceae (Evans *et al.*; 2000), as well as *Viburnum* (Adoxaceae) (Winkworth & Donoghue, 2004). The phylogenetic utility of the nuclear gene *waxy* have been demonstrated across a wide range of taxonomic levels in

different groups of plants, such as Poaceae (Yang *et al.*, 2007; Ingram and Doyle, 2003, Mason-Gamer *et al.* 1998), Malvaceae (Small, 2004), Solanaceae (Levin *et al.*, 2005; Peralta & Spooner, 2001; van der Leij *et al.*, 1991), and Convolvulaceae (Miller *et al.*, 1999). *Waxy* exhibited sufficient variability to resolve relationships including between distantly related species (e.g., wide hybridization between distantly related species within subfamily Maloideae; Evans & Campbell, 2002) and among species that gave rise to the allopolyploid *Geinae* (Colurieae: Rosaceae) (Smedmark *et al.*, 2003; Smedmark *et al.*, 2005).

Importance of a good phylogenetic framework

Madagascar is one of the eight “hottest” biodiversity hotspots based on richness and endemism of plants, and on habitat loss (estimated at >90 per cent) (Ganzhorn *et al.*, 2001). The plant diversity on Madagascar is seriously threatened: 54% of the species of Coleae are threatened with extinction (Good *et al.*, 2006)). Since conservation requires reliable information on species identity, a good phylogenetic assessment is of great importance in determining taxonomic status of organisms.

Additionally, a robust phylogenetic framework is useful in assessing the evolution of traits important to diversification (Donogue, 1989). Coleae provide a rich variety of traits including floral, fruit, and leaf characters that provide clues to the astounding recent diversification of this group.

The purpose of this study is to determine the relationships within Coleae using *waxy*, comparing and contrasting *waxy* with the existing ITS phylogeny, and evaluating the combined data phylogeny.

CHAPTER 2

MATERIALS AND METHODS

Study organisms, sampling and outgroup choice.

Materials were available for approximately 65 taxa. Two outgroup taxa were chosen: *Kigelia africana*, an Africa genus placed by Gentry (1976) in Coleeae, and *Tabebuia*, a member of Central American tribe Crescentiae that originally included Coleeae (Gentry, 1980). Outgroup taxa were chosen due to their close relationship with Coleeae (Spangler & Olmstead, 1999). Both taxa were used to root the tree in all the analyses.

DNA extraction

Total DNA was extracted from ~0.1 g of herbarium or silica gel preserved leaves. The silica gel material was collected in Madagascar between 1994 and 1998. DNA was extracted using either a modified CTAB (cetyltrimethylammonium bromide) procedure (Weigel & Glazebrook, 2002) or CTAB DNA mini-prep with 6% CTAB (Murray & Thompson, 1980; Saghai-Marooif *et al.*, 1986). In both methods, DNA was resuspended in 50 µl of distilled-deionized water.

Amplification of nDNA gene region

To amplify GBSSI, 2µl of total DNA was used in a total volume of 50µl reaction. Optimal concentrations for PCR were determined (Table 1). The first pair of primers employed were waxy13R 5'GGAGTGGCRACGTTTTTCCTT3', waxy10F 5'ACTGCTGGNGCTGATTTTATG3', and 7F 5'GYTTTSTGCATCCACAACATTGC3' (Olmstead, pers. comm.). With 10F and 13R a 650-750 bp length was expected while for 7F and 13R was of 1500 bp. The annealing

temperature was optimized for the combination of primers 10F & 13R, resulting in *Rhodocolea* amplifying at 61°C, and *Phyllarthron* and *Colea* at 60°. A hot start and touchdown procedure was employed frequently as follows: 94°C for 50 sec, 59°C for 1 min, 72°C for 1:30 min (ten cycles), 94°C for 50 sec, 56°C for 1:00 min, 72°C for 1:30 (ten cycles), 94°C for 50 sec, 54°C for 1:30 min, 72°C for 1:30 (20 cycles), and a final extension step at 72°C for 7 min. PCR products were checked on 1.2% agarose gels stained with 5µl of 1:10 ethidium bromide and viewed with ultraviolet light. Concentrations were estimated by visual comparison with bands containing known amounts of DNA.

Various Taq polymerases were used for most routine procedures. For cloning, however, *ExTaq* buffer (TaKaRa) and 0.25 µl *ExTaq* (Takara) were used to assure the 5' and 3' terminal deoxyadenina overhang. PCR were performed in an Eppendorf Thermal Cycler. When more than one product was obtained, stringency was applied by increasing the annealing temperature to yield a simple PCR product. PCR products were purified using QIAquick PCR purification kit (Qiagen). Final elution was done with buffer EB provided with the kit. Amplification products were checked on 1.2% agarose gels stained with 5µl of 1:10 ethidium bromide and viewed with ultraviolet light for presence of inserts. Concentrations were estimated by visual comparison with bands containing known amounts of DNA.

Some samples were directly sent to the department of Biochemistry and Molecular Biology at Oklahoma State University for sequencing along with downstream and upstream primers in concentrations of 10picomoles/µl. Other samples were used in cloning. Oklahoma State University used the Applied Biosystems Big Dye Terminator

1.1 cycle sequencing kit. Reactions were purified using Edge Biosystems Dye Terminator Removal System. Purified reactions were analyzed on an Applied Biosystems 3730 DNA analyzer.

Cloning

After PCR product was purified, the samples were cloned with the TOPO TA cloning kit for sequencing (Invitrogen, Carlsbad, California, USA). Ligation, transformation, and plating were carried out following the manufacturer's instructions, except that ligation and transformation volumes were halved. Ligation incubated for 30 minutes on a thermocycler at 22°C. The nutritive media for plating and growing the bacteria was Luria Bertani agar and Luria Bertani broth (Fisher, Fair Lawn, New Jersey) with 100 µg/ml of kanamycin to grow colonies with the inserted plasmid. Incubation was done in 1ml per colony and left for approximately 48 hours with vertical shaking. The DNA plasmid was purified using QiaPrep Spin Miniprep kit (Qiagen). Plasmids with inserts were screened by PCR using the M13 reverse and forward primers that come with the cloning kit.

Alignment

Sequence chromatograms were checked and edited manually in Genes Code Corporation's Sequencher, version 4.1.2 for Machintosh (Ann Arbor Michigan, USA). With cloned sequences, the vector was trimmed by imputing the flanked regions around the insert. Sometimes manual trimming was necessary. Once edited, a consensus sequence was obtained by assembling the upstream and downstream DNA sequences for each taxon. Consensus sequences were exported as text to McClade, 4.03 and further

manually adjusted. Consensus sequences were subjected to a BLAST search (Altschul *et al.*, 1997) in GenBank to test for GBSSI identity.

Primer design

From the few initial sequences obtained with the two universal pairs of primers, new primers were designed (Table 2). Once aligned, conserved regions at the beginning and end of the sequences were selected and primers were ordered from Invitrogen. Primers were designed with a similar melting point, as recommended (Strand *et al.*, 1997). Initially, four primers were designed without considering exon or intron position in order to amplify as large a portion of the gene as possible (Mort & Crawford, 2004). Primers “waxy8downs,” “waxy12ups” were designed to align with exon positions 8 and 12 of Potato *waxy* gene (Figure 1) (GenBank x58453) (van der Leij *et al.*, 1991).

Phylogenetic analyses

Phylogenetic trees were inferred using maximum parsimony (MP), and maximum likelihood (ML) optimality criterion in PAUP*4.0b10 (Swofford, 2002). In addition, data was analyzed using a Bayesian analysis approach with Mr. Bayes (v3.0; Huelsenbeck & Ronquist, 2001).

MP was set for heuristic search with “tree-bisection-reconnection” (TBR) branch swapping, collapsed branches if maximum length was zero, and all characters equally weighted and unordered. Gaps were treated as missing data. Multistate taxa were interpreted as uncertainty. A strict consensus tree was obtained. Relative measures of support for clades were estimated using Bootstrap analysis (Felsenstein, 1985), and are represented by numbers over the branches. MaxTrees was set to 2000 replicates with full heuristic search.

For both ML and Bayesian analyses, Modeltest version 3.7 (Posada & Crandall, 1999) was used in combination with PAUP* to select for the best evolutionary model for each data based on the Akaike information criterion (AIC; Akaike, 1974). ML analyses were set to a maximum of 2000 trees. TBR branch-swapping and the MulTrees options were in effect. Starting trees were obtained from the MP analysis.

Bayesian analyses were conducted with Mr. Bayes version 3.1.2. Each analysis was initiated from a random starting tree and the program was set to run four Markov chain Monte Carlo iterations (three heated and one cold) simultaneously for 50,000 generations and a tree was saved every 100 generations. The posterior probabilities for clades were estimated and are represented by numbers above the branches.

ITS and *waxy* were analyzed separately to construct phylogenetic trees: ITS alone with a total length of 720 bp (using data from Zjhra, 1998); *waxy* with a total length of 1198 bp and a subset of *waxy* that minimized missing data with 439 bp. Two combined dataset with ITS and *waxy* were analyzed: *waxy* with 1198 bp; *waxy* with 439 bp. Analyses were conducted with and without *P. cauliflorum* since it was variously placed in the *Colea* clade in the *waxy* data set. *Kigelia* and *Tabebuia* were used in all cases as outgroups. The different genera are indicated by different colors on the phylogenetic trees.

Table 1. PCR's general conditions.

Reagent	Start concentration	End concentration	Volume μ L/50 μ L
Buffer	10X	1X	5
MgCl	25mM	2.5 mM	5
dNTP	10 μ M	0.4 μ M	2
Upstream Primer	20 μ M	0.8 μ M	1
Downstream Primer	20 μ M	0.8 μ M	1
BSA	100%		1
Water			to volume

Table 2. Primers used in PCR amplification and sequencing of *waxy*.

Pair of primers	Primer sequence (with forward and reverse)	Sequence length product expected (bp.)
WaxyFColeae WaxyRColeae	5'gccatgccaccaggtcacc 3' 5'tggtctcatccaattgcatgccatgcg 3'	650
Coleaeupstream1 Coleaedownstream1	5'catgccaccaggtcacc 3' 5'ccaattgcatgccatgcgatac 3'	No product
Coleaeupstream2 Coleaedownstream2	5' ctggtgaatcaggtacatctg 3' 5'gttctcaatgactacctccaagagatc3'	520
GBSS1coleaeandredown GBSS1coleaeandreups	5' gtgagatagccatgagttgaag 3' 5' gatacggaacagtaagagccctag 3''	800
Waxy8downs waxy12ups	5'ctggatgmaggctggaattrtaga atc 3' 5'gttgaagcacagataggtacc 3'	780




Figure 2. Diagram of granule-bound synthase gene from *Solanum tuberosum*. Arrows indicate locations and directions of waxy8ups and waxy12downs. (Diagram adapted from van der Leij *et al.*, 1991)

CHAPTER 3

RESULTS

Amplification and alignment:

A total of 24 ingroup taxa were successfully amplified (Fig. 2) which represent approximately 36% of the total species described, and encompasses three of the four genera. Despite intensive efforts, *Ophiocolea* failed to amplify. The alignment of the sequences of *Colea*, *Rhodocolea*, and *Phyllarthron* were straightforward. Outgroup alignment of *Tabebuia* and *Kigelia* was straight forward up to exon 12, after which variability made the alignment ambiguous. This last ambiguous portion was eliminated from the analysis; this variability was potentially a result of low sequence signal.

Phylogenetic analysis.

Maximum Parsimony, Maximum Likelihood, and Bayesian analyses were performed on 3' portion of *waxy* obtained in this study. The maximum bp length encompassed 1198 bp (“*waxy*” data set) from exon 8. Missing data resulted from using different combinations of primers for successful amplification. To explore the impact of missing data, a data subset that minimized missing data was also analyzed. This data set was 439 bp long (“subset of *waxy*”). Previous analysis with *waxy* under MP showed *P. cauliflorum* as a member of *Colea* clade. To test the impact of *P. cauliflorum* on the analysis, analyses were conducted with and without it. Trees obtained with *P. cauliflorum* are not show since placement of the taxa was uncertain.

MP vs. ML vs. bayesian analysis.

ITS: this data set included 21 ingroup taxa and 2 outgroup. The ITS sequences were 720 bp in length that included ITS 1, 5.8 s and ITS 2 from nuclear ribosomal DNA

and included 106 (11.08%) parsimony-informative character-sites. The score of the best tree found was $L = 286$ and 23 trees were retained from which a strict consensus tree was obtained (Fig. 3A). In this tree, two subclades of *Phyllarthron* showed up: *Phyllarthron* I is sister to *Phyllarthron* II and sister to the rest. Taxa that belong to different populations but the same species came out as more closely related except for *P. madagascariense* and *R. nobilis*. Support was over 70% for most of the branches, except *Colea* and some branches in *Phyllarthron* I. The ML tree (Fig 3B) is more resolved than MP, showing *C. cava* as sister of the rest of *Colea*, a monophyletic *Rhodocolea*, and two clades of *Phyllarthron*. The best tree score was $-\ln L = 2606.09118$. Topology of the Bayesian analyses tree (Fig.3C) is similar to the one obtained by Maximum likelihood analysis, with a minor difference in the placement of the two accessions of *Rhodocolea nobilis*.

Subset of *waxy*: No missing data was present except for *Kigelia* and *Tabebuia*. Sixty-one characters (13.8 %) were parsimony-informative. The best tree score was = 219. Two trees were retained and a strict consensus obtained (Fig. 4A). As in ITS, *Colea* and *Rhodocolea* were monophyletic under ML and Bayesian analyses (Fig. 4B, C). However, *waxy* results in monophyletic *Colea* and *Rhodocolea* under MP (Fig. 4A) whereas ITS does not. The best tree score was $-\ln L = 1759.15410$. Bayesian tree (Fig. 4C) has the same topology as the ML tree.


3' *waxy*: A total of 23 ingroup taxa were analyzed for *waxy* with 1198 characters, from which 196 were parsimony-informative. Score of best tree was 391 and number of trees retained was equal to 45. The topology of the MP tree is similar to the one obtained from the subset data analysis (Fig. 5A), differing only in slight variations in the relationships within *Phyllarthron* II. Branch support is higher for 3' *waxy*, in general,

than for subset *waxy*. Topology is consistent among the three analyses of the 3' *waxy* (Fig. 5A, B, C). The best tree score found was $-\ln L$ 3807.66189 for ML.


Combined data 1: This data set included 20 taxa with both ITS and *waxy* sequences, using 3' *waxy* (1198 bp). A total of 18 ingroup taxa were included. Out of 1907 characters, there were 161 parsimony-informative characters. The best tree found was= 636 and number of trees retained was= 10. The combined data analyses trees (Fig. 6A, B, C) are similar to separate analyses, but with the two *Phyllarthron* clades adjacent and *Rhodocolea* within *Colea*. ML tree score for best tree found was $-\ln L$ 6173.42045.

Combined data set 2: This dataset included the subset of *waxy* plus ITS for the 18 included ingroup taxa. There were 125 informative characters. This data set included 441 bp length from *waxy* and 720 of ITS data. The best parsimony tree was=475. The number of trees retained was=31. Results are similar to the combined data set 1, but with more resolution within clades and monophyly of *Colea* in the Bayesian tree (Fig. 7A, B, C). The best tree score found for ML analysis was $-\ln L$ 4219.36415.


Congruence between individual gene trees vs. combined data: Tendencies are mostly the same when data is combined in comparison with individual data (Fig. 8). However, in the combined data, *Colea* is paraphyletic.




C. cava




C. gentry



C. muricata




C. ramiflora




C. systmae


A) *Colea*




P. articulatum



P. bernieranum




P. madagascariense



P. nocturnum



P. sahamalenze



P. vokoanensis

B) *Phyllarthron*




Figure 2. Map of distribution of Coleae's species. Shaded areas: 2000 forest cover; dark green=rain forest, yellow= spiny desert. Pale green= mangrove forest, blue= cloud cover. Blue dots= species localities for A) *Colea*, B) *Phyllarthron*, and C) *Rhodocolea*. (Maps by Dr. Good)




Figure 3. Phylogenetic trees from analyses of ITS. A) Strict consensus tree of 23 trees (L=286) from MP analyses. Bootstrap values are numbers above the branches. B) ML tree (-ln L 2606.09118). C) Bayesian tree showing posterior clade probabilities.




Figure 4. Phylogenetic trees from analysis of a subset of *waxy*. A) Strict consensus tree of 2 trees (L=219) from MP analysis. Bootstrap values are numbers above the branches. B) ML tree (-ln L1759.15410). C) Bayesian tree showing posterior clade probabilities.




Figure 5. Phylogenetic trees from analysis of *waxy* (1198 bp). A) Strict consensus tree of 45 trees (L=391) from MP analyses. Bootstrap values are numbers above the branches B) ML tree (-ln L 3807.66189). C) Bayesian tree showing posterior clade probabilities.




Figure 6. Phylogenetic trees from analysis of combined data 1 (ITS+waxy). A) Strict consensus tree of 10 trees (L= 636) from MP analyses. Bootstrap values are numbers above the branches. B) ML tree (-ln L 6173.42045). C) Bayesian tree showing posterior clade probabilities.




Figure 7. Phylogenetic trees from analysis of combined data 2 (ITS+subset *waxy*). A) Strict consensus tree of 31 trees (L=475) from MP analyses of the combined data 2. Bootstrap values are numbers above the branches. B) ML tree (-ln L 4219.3615). C) Bayesian tree showing posterior clades probabilities.




Figure 8. Bayesian phylogenetic trees of A) ITS, B) *waxy*, C) combined data, and D) phylogenetic representation of Gentry's (1980) taxonomic relationships within Coleeae.

Table 3. Summarized results of modeltest:

Data set	Model selected	Substitution model	Bases frequencies	Proportion of invariable sites	Gamma distribution shape
ITS	GTR+I+G	R (a) [A-C] = 1.1082 R (b) [A-G] = 1.6632 R(c) [A-T] = 0.9372 R (d) [C-G] = 0.5916 R (e) [C-T] = 4.2120 R (f) [G-T] = 1.0000	A= 0.2369; C= 0.3059; T= 0.1791 G= 0.2782.	0.3700	0.7983.
Subset de <i>waxy</i>	TVM+G	R(a) [A-C] = 1.2551, R(b) [A-G] = 3.1914, R(c) [A-T] = 0.8781, R(d) [C-G] = 2.1382, R(e) [C-T] = 3.1914, R(f) [G-T] = 1.0000.	A= 0.3031 C= 0.1325 T= 0.3615 G= 0.2030	0	0.9577
<i>Waxy</i>	TrN+G	:R(a) [A-C] = 1.0000, R(b) [A-G] = 2.6295, R(c) [A-T] = 1.0000, R(d) [C-G] = 1.0000, R(e) [C-T] = 3.8432, R(f) [G-T] = 1.0000.	A= 0.2953, C= 0.1600, T= 0.3282 G= 0.2165	0	0.4761
Combined data 1	TrN+G	R(a) [A-C] = 1.0000, R(b) [A-G] = 2.4952, R(c) [A-T] = 1.0000, R(d) [C-G] = 1.0000, R(e) [C-T] = 4.0940, R(f) [G-T] = 1.0000.	A= 0.2691, C= 0.2183, T= 0.2733, G= 0.2392	0.3774	0.8258
Combined data 2	TrNeF+I+G	R(a) [A-C]= 1.0000, R(b) [A-G] = 2.4787, R(c) [A-T] = 1.0000, R(d) [C-G] = 1.0000, R(e) [C-T] = 3.5945, R(f) [G-T] = 1.0000.	Equal frequencies	0.3338	0.7905. □

CHAPTER 4

DISCUSSION

Effect of sampling: differences in resolution among the main clades as well as within clades probably derived from the differences in both the number and species sampled in ITS and *waxy* data sets. Complete absence of members of genus *Ophiocolea* may also account for the relationships observed. In Zjhra's (2003) analysis of ITS with 65 taxa, *Colea* was sister to *Ophiocolea*. In the larger dataset of ITS, *Phyllarthron* is also paraphyletic. This can be due to the paucity of sampling within *Phyllarthron* itself, since phylogenies constructed with chloroplast DNA including broad representation of *Phyllarthron* taxa showed *Phyllarthron* as monophyletic (Zjhra, 2004).

Effect of missing data: When analyzing *waxy* data with a subset or the total length of the data set, better resolution for most of the branches came out with the total length of the sequences than for a subset, but phylogenetic relationships stayed the same for most of the branches. Cracraft and Helm-Bychowski (1991) analyzed primate relationships using mitochondrial DNA and found that different subsets of the data resolved consistently with few exceptions, and although the tendency is to improve the resolution with the addition of data, this is not always true. This was true for the Coleae *waxy* data, where the analyses of a subset showed better resolution for *Colea*, whereas within *Phyllarthron* the resolution improved with adding more data. This can be due to few phylogenetically informative characters in the original short *Colea* sequences. Missing data can be an obstacle when reconstructing phylogenetic relationships (Donogue *et al.*, 1989), leading to multiple shortest trees and poorly resolved consensus trees (Gauthier, 1986). Although missing data can result in inaccurate phylogenies, the

mechanisms still remain unclear. Huelsenbeck (1991), for instance, proposed that this is due to the percentage of equivocally resolved ancestral characters. By computer simulations, Wiens (2003) found support for the hypothesis that the lack of resolution when using incomplete taxa is primarily one of including too few characters rather than including too many missing data cells.

Utility of *waxy* vs. ITS. The internal transcriber spacer of the nuclear rDNA has been used to study relationships among closely related species, but frequently is not variable enough to distinguish species in some groups (Mason-Gamer *et al.*, 1998). In *Hibiscus* Sect. *Muenchhusia* (Small, 2004) ITS data contained 17 nucleotide substitutions out of a total of 683 bp aligned nucleotides, with only 2 potentially phylogenetically informative characters. GBSSI, on the other hand, consisted of a total of 1972 bp with 28 phylogenetically informative. However, in other cases, ITS have shown more resolving power than *waxy*: in *Ipomoea* (Miller *et al.* 1999) a total length of 573 nucleotides had 163 phylogenetically informative sites, contrasting with *waxy* which showed just 86 potentially informative sites in a final alignment of 651 bp length. In Coleae, ITS shows enough variability to resolve relationships among species, even with fewer taxa. *Waxy* was also useful in resolving phylogenetic relationships within species, even with a very short subset of data. The informativeness of ITS (14.68%) and *waxy* (13.89%) was equivalent. However, GBSSI is a longer region and therefore potentially provides a greater number of phylogenetically informative sites.

Common terminal branch relationships among ITS data and *waxy* include *P. vokoanina* 709 and *P. madagascariense* CapEst (*P. madagascariense* 967 in *waxy*), which is also supported by previous analyses of ITS (Zjhra, 2003). Finally, monophyly of

Colea and *Rhodocolea* are supported by both ITS and *waxy* as well as the presence of two subclades of *Phyllarthron*.

Combined data: In general, there was less resolution in the combined data sets compared to the individual data sets. This can be due to differences in sampling between the *waxy* and ITS data sets. Only in taxa better represented, such as *Phyllarthron*, the resolution is consistent between the combined and separate data analyses. The combined data sets showed *Colea* as paraphyletic and *Rhodocolea* unresolved. Also, in both combined data sets, *Colea* and *Rhodocolea* came out as a single clade. This finding conflict with results derived from three cpDNA data sets where *Rhodocolea* is sister to the rest of the tribe (Zjhra *et al.* 2004). When taxa don't have strong support, they can appear anywhere in the tree; *Colea* lacked strong support in both of the individual data analyses.

Phylogeny methods: Bayesian analysis generally resulted in better resolution than ML and MP. Posterior probabilities tend to be higher in all the cases than bootstrap values. However, Bayesian posterior probabilities determine the strength of the data in supporting particular nodes, whereas bootstrap values indicate areas where additional data is needed to resolve relationships (Miller *et al.* 2004).

Phylogenetic relationships: The main phylogenetic relationships observed are monophyly of *Colea* and *Rhodocolea*, and two subclades of *Phyllarthron*. These relationships are also supported in the broader phylogenetic analysis of ITS (Zjhra, 2003) with 61 ingroup taxa. When comparing phylogenetic analyses of ITS data with *waxy* data, the major differences are that *Rhodocolea* is sister to *Phyllarthron* 1 which is sister of *Colea* and *Phyllarthron* 2 (ITS data), whereas *Phyllarthron* 2 is sister to a

monophyletic *Colea* and *Rhodocolea (waxy)*. In the 61 taxa ITS phylogeny (Zjhra, 2003), *Rhodocolea* was sister of the rest of the tribe. The results from *waxy* place *Rhodocolea* in a derived position, whereas Gentry (1988) placed *Rhodocolea* in a primitive position within the tribe (Figure 8).

Finally, phylogenetic relationships were surprisingly robust to the effects of missing data, with branch support the major difference between analyses of the entire vs. subset of *waxy*. ITS shows enough variability to resolve relationships among species, even with fewer taxa. *Waxy* was also useful in resolving phylogenetic relationships within species, even with a very short subset of data. Monophyly of *Colea* and *Rhodocolea* are supported by both ITS and *waxy* as well as the presence of two subclades of *Phyllarthron*. Less resolution in most of the branches can be observed in combined data sets when compared with individual data.

LITERATURE CITED

- Álvarez, I; & Wendel, J. F. 2003. Ribosomal ITS sequences and plant phylogenetic inference. *Molecular Phylogenetics and Evolution*, 29: pp. 417-434.
- Akaike, H. 1974. A new look at statistical model identification. *IEEE. Trans. Automatic Control* 19: 716-723.
- Altschul, S. F; Madden, T. L; Schäffer, A. A; Zhang, J; Zhang, Z; Miller, W; & Lipman, D. J. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search program. *Nucleic Acids Research*, 25: pp. 3389-3402.
- APG II (Angiosperm Phylogeny Group II). 2003. An update of the Angiosperms Phylogeny Group classification for the orders and families of flowering plants: APG II. *Botanical Journal of the Linnean Society*. 141: pp. 399-436.
- Baillon, H. 1887. Notes sur les Crescentiées. *Bull. Soc. Linn. Paris*. 1: 678, 680, 683, 688, 690, 695.
- Baillon, H. 1888. *Histoire des plantes* 10. Hachette, Paris.
- Baldwin, B. G. 1992. Phylogenetic utility of the internal transcribed spacers of nuclear ribosomal DNA in plants: an example from the Compositae. *Molecular Phylogenetics and Evolution*, 29: pp. 262-265.
- Bojer, W. 1837. *Hortus Mauritianus*. Aime Mararot & Co., Mauritius.
- Candole, A. P. de. 1838. *Revue sommaire de la famille des Bignoniacees*. *Bibliothèque Universelle de Geneve*, Geneve.
- Candole, A. P. de. 1845. *Prodomus systematis naturalis regni vegetabilis* 9. Treuttel et Würtz, Paris.
- Cracraft, J; & Helm-Bychowski, K. 1991. Parsimony and phylogenetic inference using DNA sequences: some methodological strategies. In: Miyamoto MM, Cracraft, J eds. *Phylogenetic analysis of DNA sequences*. New York: Oxford University Press: pp. 184-220.
- Cronn, R., Small, R., Haselkorn, T. & Wendel, J. 2003. Cryptic repeated genomic recombination during speciation in *Gossypium gossypioides*. *Evolution* 57: pp. 2475-2489.
- De Jussieu, A. 1789. *Genera Plantarum, secundum ordines naturales disposita juxta methodum in Horto Regio Parisiensi exaratam*. MS. notes. Parisiis, 1789.
- Denyer, K., Johnson, P., Zeeman, S., & Smith, A. 2001. The control of amylose synthesis. *Journal of Plant Physiology*. 158: pp. 479-487.
- Donoghue, M. J. 1989. Phylogenies and the analysis of evolutionary sequences, with examples from seed plants. *Evolution*, 43: pp. 1137-1156.
- Donoghue, M. J; Doyle, J. A; Gauthier, J; Kluge, A. G; & Rowe, T. 1989. The importance of fossils in phylogeny reconstruction. *Annu. Rev. Ecol. Syst.*, 20: pp. 431-460.

- Doyle, J. J. 1992. Gene trees and species trees: molecular systematics as one-character taxonomy. *Systematic Botany*, 17(1): pp. 144-163.
- Echt, G. S. & Schwartz, D. 1981. Evidence for the inclusion of controlling elements within the structural gene *waxy* locus in maize. *Genetics*, 99: pp. 275-284.
- Evans, R. C.; Alice, L. A.; Campbell, C. S.; Kellog, E, A. & Dickinson T. A. 2000. The granule-bound starch synthase (GBSS1) gene in the Rosaceae: Multiple loci and phylogenetic utility. *Molecular Phylogenetics and Evolution*, 17(3): pp. 388-400.
- Evans, R. C; & Campbell, C. S; 2003. The origin of the apple subfamily (Maloideae: Rosaceae) is clarified by DNA sequence data from duplicated GBSSI genes. *American Journal of Botany*, 89(0): pp. 1478-1484.
- Felsenstein, J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution*, 39: pp. 783-791.
- Ganzhorn, J. U; Lowry II, P. P; Schatz, G. E; & Sommer S. 2001. Short Communication. The biodiversity of Madagascar: one of the world's hottest hotspots on its way out. *Oryx*, 35 (4): pp. 346-348.
- Gauthier, J. 1986. Saurischian monophyly and the origin of birds. *Mem. Calif. Acad. Sci.*, 8:Pp. 1-47.
- Gautier, L. & Goodman, S. M. 2003. Introduction to the Flora of Madagascar. S. Goodman and J. Benstead, eds. Natural History of Madagascar. University of Chicago Press. Pp. 229-250.
- Gentry, A, H. 1976. Relationship of the Madagascar Bignoniaceae: A striking case of convergent evolution. *Plant Syst. Evol.* 126: pp. 255-266.
- Gentry, A. H. 1980. Bignoniaceae. Part 1. Tribes Crescentiae and Tourrentiae. *Flora Neotropica Monograph* 25: pp. 1-130.
- Gentry, A. H. 1988. Distribution and evolution of the Madagascar Bignoniaceae. *Monogr. Syst. Bot. Missouri Bot. Gard.* 25: pp. 175-185.
- Giles, E. 1988. L'equilibre des ecosystems forestiers a Madagascar actes d'um seminaire international. IUCN. Pp.155-171.
- Good, T, Zjhra, M. & Kremen, C. 2006. Addressing data deficiency in classifying extinction risk: a case study of a radiation of Bignoniaceae from Madagascar. *Conservation Biology*, 20(4): pp. 1099-1110.
- Ingram, A; & Doyle, J. 2003. The origin and evolution of *Eragrostis tef* (Poaceae) and related polyploids: evidence from nuclear gene *waxy* and plastid *rps16*. *American Journal of Botany*. 90 (1): pp. 116-122.
- Jury, M. R. 2003. The climate of Madagascar. S. Goodman and J. Benstead, eds. Natural History of Madagascar. University of Chicago Press. Pp. 75-87.
- Levin, R. A.; Watson, K. & Bohs, L. 2005. A four-gene study of evolutionary relationship in *Solanum* section *Acanthophora*. *American Journal of Botany* 92(4): pp. 603-612.

- Mason-Gamer, R. J; Weil, C. F; & Kellogg, A. 1998. Granule-bound starch synthase: structure, function, and phylogenetic utility. *Molecular Biology and Evolution*. 15: pp. 1658-1673.
- Merida, A. Rodriguez-Galan, J. M. & Romero, J. 1991. Expression of the granule-Bound Starch Synthase I (Waxy) Gene from Snapdragon is developmentally and Circadian Clock Regulated. *Plant Physiology*, 120: pp. 401-409.
- Miller, R. E, & Rausher, M. D. Manos, P. S. 1999. Phylogenetic Systematics of Ipomoea (Convolvulaceae) Based on ITS and *Waxy* sequences. *Systematic Botany*, 24(2):pp. 209-227.
- Mort, M. E. & Crawford D. J. 2004. The continuing search: low-copy nuclear sequences for lower level plant molecular phylogenetic studies. *Taxon*, 53(2): pp. 257-261.
- Mort, M. E; Archibald, J. K; Randle, C. P; Levsen, N. D; O'Leary, T. R; Topalov, K. Wiegand, C. M. & Crawford, D. J. 2007. Inferring phylogeny at low levels: Utility of rapidly evolving cpDNA and nuclear ITS. *American Journal of Botany*, 94(2): pp. 173-183.
- Murray, M. G; & Thompson, W. F. 1980. Rapid isolation of high molecular weight plant DNA. *Nucleic Acids Res.* 8: pp. 4321-4325.
- Nixon, K. C; & Carpenter. 1996. On consensus, collapsibility, and clade concordance. *Cladistics* 12: pp. 305-321.
- Olmstead, R.G; Bremen, B; Scout, K. M; Palmer & J. D. 1993. A parsimony analysis of the Asteridae sensu lato based on *rbcL* sequences. *Ann. Missouri Bot. Gard.*, 80: pp. 700-722.
- Perrier de la Bâthie, H. 1938a. *Les Bignoniacées* de la region Malgache. *Ann. Mus. Col. Marseille*, 46: pp. 1-101.
- Perrier de la Bâthie, H. 1938b. *Bignoniacées*. *Flore de madagascar*, 178: pp. 1-91. Muséum National d'Histoire Naturelle, Paris.
- Peralta, I. & Spooner, D. 2001. Granule-Bound starch synthase (GBSS1) gene phylogeny of wild tomatoes (*Solanum* L. section *Lycopersicon* [Mill.] Wettst. Subsection *Lycopersicon*). *American Journal of Botany*, 88(10): pp. 188-1902.
- Posada, D. & Crandall, KA. 1999. Modeltest: testing the model of DNA substitution. *Bioinformatics*, 14 (9): pp. 817-818.
- Rabinowitz P. D.; Coffin M. F.; Falvey D. 1982. Salt diapers bordering the continental margin of northern Kenya and Southern Somalia. *Science*, 215: pp. 663-664.
- Rabinowitz P. D.; Coffin M. F.; Falvey D. 1983. The separation of Madagascar and Africa. *Science*, 220: pp. 67-69.
- Saghai- Maroof, K. M; Soliman, R. A; Jorgensen, A. A; & Allard, R. W. 1984. Ribosomal DNA spacer-length polymorphism in barley: Mendelian inheritance, chromosomal location, and population dynamics. *Proc. Natl. Acad. Sci., USA*. 81: pp. 8014-8018.

- Sang, T. 2002. Utility of low-copy nuclear gene sequences in plant phylogenetics. *Critical Reviews in Biochemistry and Molecular Biology*, 37(3): pp. 121-147.
- Seemann, B. 1860. Synopsis Crescentiacearum: an enumeration of all the crescentaceous plants at present known. *Trans. Linn. Soc. London*. 23: pp. 1-22.
- Small, R. L. 2004. Phylogeny of *Hibiscus* sect. *Muenchhusia* (Malvaceae) based on chloroplast *rpL16* and *ndhF*, and nuclear ITS and GBSS1 sequences. *Systematic Botany*, 29(2): pp. 385-392.
- Smedmark, J. E. E., Eriksson, T. Evans, R. C. & Campbell C. S. 2003. Ancient allopolyploid speciation in Geinae (Rosaceae): evidence from nuclear granule-bound starch synthase (GBSSI) gene sequences. *Syst. Biol.* 52: pp. 374-385.
- Smedmark, J. E; Eriksson, T; Bremer, B. 2005. Allopolyploid evolution in Geineae (Colurieae: Rosaceae) – building reticulate species trees from bifurcating gene trees. *Organisms, Diversity and Evolution*, 5: pp. 275-283.
- Stebbins, G. L. 1974. Flowering plant: evolution above the species level. Belknap Press, Cambridge, MA.
- Storey, M; Mahoney, J. J; Saunders, A. D. Duncan, R. A; Kelley, S. P; & Coffin, M. F. 1995. Timing of hot spot-related volcanism and the breakup of Madagascar and India. *Science*, 267: pp. 852-855.
- Strand, A. E; Leebens-Mack, J. & Milligan, G.B. 1997. Nuclear DNA-based markers for plant evolutionary biology. *Molecular Ecology*, 6: pp. 113-118.
- Swofford, D. L. 2002. PAUP*. Phylogenetic analysis using parsimony (and other methods). Version 4.0b10. Sunderland: Sinauer Associates.
- van der Leij, F. R., Visser, R. G, Ponstein, A. S., Jacobsen, E. & Feenstra, W. 1991. Sequence of the structural gene for granule-bound starch synthase of potato (*Solanum tuberosum* L.) and evidence for a single point deletion in the *amf* allele. *Mol. Gen. Genet.*, 228: pp. 240-248.
- Walters, H. 1971. Ecology of tropical and subtropical vegetation. Oliver & Boyd, Edinburgh.
- Weigel, D; & Glazebrook, J. 2002. Arabidopsis: a laboratory manual. Cold Spring Harbor laboratory press. Cold Spring Harbor, New York. Pp. 165-166. *Syst.*, 52(4): pp. 528-538.
- Wiens, J. 2003. Missing data, incomplete taxa, and phylogenetic accuracy. 2003. *Syst. Biol.*, 52(4): pp. 528-538.
- Winkworth, R. & Donoghue, M. 2004. *Viburnum* phylogeny: evidence from the duplicated nuclear gene GBSS1. *Molecular Phylogenetics and Evolution*, 33: pp. 109-126.
- Yang, H. Q.; Peng, S. & Li, D. Z. 2007. Generic delimitations of *Schizostachyum* and its allies (Gramineae: Bambusoideae) inferred from GBSSI and *trnL-F* sequence phylogenies. *Taxon*. 56(1): pp. 45-54.

- Zjhra, M.; Sytsma, K.; Olmstead, R. 2004. Delimitation of Malagasy Coleeae and implications for fruit evolution in Bignoniaceae inferred from a chloroplast DNA phylogeny. *Plan. Syst. Evol.* 245: pp. 55-67.
- Zjhra, M.L. 2003. Malagasy Bignoniaceae: phylogenetics and evolution of pollination systems. S. Goodman and J. Benstead, eds. Natural History of Madagascar. University of Chicago Press. Pp. 425-430.
- Zjhra, M. L. 2006. New taxa of Coleeae from Madagascar I. A collection from Masoala peninsula. *Ann. Bot. Fennici*, 43: pp. 225-239.