

References

- [1] Albertson, M., Pach, J., Young, M., Disjoint homometric sets in graphs. *Ars Mathematica Contemporanea*, 4(1), 2011. <https://doi.org/10.26493/1855-3974.174.027>
- [2] Alon, N., Problems and results in Extremal Combinatorics-III. *Journal of Combinatorics*, 7(2-3):233–256, 2016.
- [3] Axenovich, M. and Özkahya, L., On homometric sets in graphs. *Electronic Notes in Discrete Mathematics*, 38: 83–86, 2011.
- [4] Bollobás, B., Kittipassorn, T., Narayanan, B., Scott, A., Disjoint induced subgraphs of the same order and size. *European Journal of Combinatorics*, 49:153–166, 2015. <https://doi.org/10.1016/j.ejc.2015.03.005>
- [5] Caro, Y. and Yuster, R., Large disjoint subgraphs with the same order and size, *European Journal of Combinatorics*, 30, no. 4, 813–821, 2009. <https://doi.org/10.1016/j.ejc.2008.09.001>
- [6] Fulek, R. and Mitrović, S., Homometric sets in trees. *European Journal of Combinatorics*, 35:256–263, 2014.
- [7] Lovász, L., Kneser’s conjecture, chromatic number, and homotopy, *Journal of Combinatorial Theory, Series A*, 25, no. 3, 319–324, 1978. [https://doi.org/10.1016/0097-3165\(78\)90022-5](https://doi.org/10.1016/0097-3165(78)90022-5)