

References

- [1] M. Albertson, Turán theorems with repeated degrees, *Discrete Mathematics*, 100 (1992) 235–241. [https://doi.org/10.1016/0012-365X\(92\)90644-U](https://doi.org/10.1016/0012-365X(92)90644-U)
- [2] M. Albertson, People who know people, *Mathematics Magazine*, 67 (1994) 278–281. <https://doi.org/10.2307/2690847>
- [3] M. Albertson, The irregularity of a graph, *Ars Combinatoria*, 46 (1997) 219–225.
- [4] M. Albertson, D. Berman, Ramsey graphs without repeated degrees, *Congressus Numerantium*, 83 (1991) 91–96.
- [5] M. Albertson, D. L. Boutin, Lower bounds for constant degree independent sets, *Discrete Mathematics*, 127 (1994) 15–21. [https://doi.org/10.1016/0012-365X\(92\)00463-2](https://doi.org/10.1016/0012-365X(92)00463-2)
- [6] T. Biedl, D. F. Wilkinson, Bounded-degree independent sets in planar graphs, *Theory of Computing Systems*, 38 (2005) 253–278. <https://doi.org/10.1007/s00224-005-1139-0>
- [7] B. Bollobás, P. Erdős, M. Simonovits, On the structure of edge graphs II., *Journal of the London Mathematical Society*, 12 (1976) 219–224. <https://doi.org/10.1112/jlms/s2-12.2.219>
- [8] P. Borg, Y. Caro, K. Fenech, Irregular independence and irregular domination, *Discrete Applied Mathematics*, in press, <https://doi.org/10.1016/j.dam.2018.07.030>
- [9] L. Caccetta, P. Erdős, K. Vijayan, Graphs with unavoidable subgraphs with large degrees, *Journal of Graph Theory*, 12 (1988) 17–27. <https://doi.org/10.1002/jgt.3190120104>
- [10] Y. Caro, A. Hansberg, M. Henning, Fair domination in graphs, *Discrete Mathematics*, 312 (2012) 2905–2914. <https://doi.org/10.1016/j.disc.2012.05.006>
- [11] Y. Caro, A. Hansberg, R. Pepper, Regular independent sets, *Discrete Applied Mathematics*, 203 (2016) 35–46. <https://doi.org/10.1016/j.dam.2015.09.020>
- [12] Y. Caro, R. Yuster, C. Zarb, Ramsey numbers for degree-monotone paths, *Discrete Mathematics*, 340 (2017) 124–131. <https://doi.org/10.1016/j.disc.2016.07.025>
- [13] G. Chartrand, M. S. Jacobson, J. Lehel, O. R. Oellerman, S. Ruiz, F. Saba, Irregular networks, *Congressus Numerantium*, 64 (1988) 187–192.
- [14] G. Chen, R. Schelp, Ramsey problems with bounded degree spread. *Combinatorics, Probability and Computing*, 2 (1993) 263–269. <https://doi.org/10.1017/S0963548300000663>
- [15] V. Chvátal, V. Rödl, E. Szemerédi, W. T. Trotter, The Ramsey number of a graph with bounded maximum degree, *Journal of Combinatorial Theory (B)*, 34 (1983) 239–243. [https://doi.org/10.1016/0095-8956\(83\)90037-0](https://doi.org/10.1016/0095-8956(83)90037-0)
- [16] V. Chvátal, E. Szemerédi, On the Erdős–Stone theorem, *Journal of the London Mathematical Society (2)*, 23 (1981) 207–214. <https://doi.org/10.1112/jlms/s2-23.2.207>

- [17] P. Erdős, G. Chen, C. Rousseau, R. Schelp, Ramsey problems involving degrees in edge-colored complete graphs of vertices belonging to monochromatic subgraphs, *European Journal of Combinatorics*, 14 (1993) 183–189. <https://doi.org/10.1006/eujc.1993.1023>
- [18] P. Erdős, A. H. Stone, On the structure of linear graphs, *Bulletin of the American Mathematical Society*, 52 (1946) 1089–1091.
- [19] S. P. Radziszowski, Small Ramsey numbers, *Electronic Journal of Combinatorics*, DS1.15 (2017), 104 pages.
- [20] Ľ. Šolteš , Nonbipartite graphs with the repeated degree property, *Journal of Graph Theory*, 26 (1997) 17–25. [https://doi.org/10.1002/\(SICI\)1097-0118\(199709\)26:1<17::AID-JGT3>3.0.CO;2-K](https://doi.org/10.1002/(SICI)1097-0118(199709)26:1<17::AID-JGT3>3.0.CO;2-K)
- [21] Zs. Tuza, How to make a random graph irregular, *Random Structures and Algorithms*, 6 (1995) 323–329. <https://doi.org/10.1002/rsa.3240060219>